

11. Specielle behandlings scenarier

HIV-positive patienter med germinative tumorer skal behandles svarende til patienter som ikke har denne infektion. Aktiv antiviral behandling kan gives sideløbende med kemoterapi.

Patienter med **udbredte lungemetastaser** (hvide lunger) kan opstartes i reduceret dosis (2-3 dages cisplatin og etoposid). Patienter med begyndende **organsvigt** kan opstarte behandling umiddelbart og afvente orchiectomi til efter afsluttet kemoterapi. Hos patienter med **nyresvigt** skal behandling med bleomycin undgås.

Referencer

1. Powles T, Robinson D, Stebbing J et al. Highly active antiretroviral therapy and the incidence of non-AIDS-defining cancers in people with HIV infection. *J Clin Oncol* 2009;27:884-890.
2. Massard C, Plantade A, Gross-Goupil M, et al. Poor prognosis nonseminomatous germ-cell tumours (NSGCTs): should chemotherapy doses be reduced at first cycle to prevent acute respiratory distress syndrome in patients with multiple lung metastases? *Ann Oncol* 2010;21:1585-1588.
3. Collette L, Sylvester RJ, Stenning SP et al. Impact of the treating institution on survival of patients with "poor-prognosis" metastatic nonseminoma. European Organization for Research and Treatment of Cancer Genito-Urinary Tract Cancer Collaborative Group and the Medical Research Council Testicular Cancer Working Party. *J Natl Cancer Inst* 1999;91:839-846.